

# Cobweb posets as noncommutative prefabs

A. Krzysztof Kwaśniewski

High School of Mathematics and Applied Informatics  
Kamienna 17, PL-15-021 Białystok, Poland

## Summary

A class of new type graded infinite posets with minimal element is introduced. These so called cobweb posets proposed recently by the present author constitute a wide range of new noncommutative and nonassociative prefab combinatorial schemes' examples with characteristic graded sub-posets as primes. These schemes are defined here via relaxing commutativity and associativity requirements imposed on the composition in prefabs by the fathers of this fertile concept. The construction and the very first basic properties of cobweb prefabs are disclosed. An another new type prefab example with single valued commutative and associative composition is provided. "En passant" though not by accident - we discover new combinatorial interpretation of all classical  $F$  - *nomial* coefficients hence specifically incidence coefficients of reduced incidence algebras of full binomial type are given a new cobweb combinatorial interpretation also.

AMS Classification Numbers: 05C20, 11C08, 17B56 .

## 1 Introduction

The concept of prefab (with associative and commutative composition) was introduced in [1], see also [2,3]. Here we shall deliver a class of similar combinatorial structure of new type based on the so called cobweb posets. For the sake of completeness we recall in Section 1. the definition of a cobweb poset as well as a combinatorial interpretation of its characteristic binomial-type coefficients (for example- fibonomial ones) [4,5]. In Section 2. after relaxing associativity and commutativity requirements imposed on the composition in prefabs by the authors of this concept [1] we observe that the vast family of all cobweb posets becomes by construction a new type of *nonassociative* non-commutative prefabs' subclass. The very first basic properties of these cobweb prefabs are shown up. As a result a class of new type of graded infinite posets with minimal element are employed here as an enveloping framework for the completely new class of combinatorial prefab structures with noncommutative and *nonassociative* composition (synthesis) of its objects since now on called **prefabiants**. Cobweb infinite posets  $P$  are designated uniquely by any cobweb admissible sequence of integers  $F = \{n_F\}_{n \geq 0}$  and are by construction endowed with self-similarity property. Namely at each graded level vertex a family of infinite cobweb sub-posets isomorphic to  $P$  may be rooted. The number of finite characteristic sub-posets (prime *prefabiants*) of  $P$  in between levels of graded Hasse digraph of  $P$  is given by  $F$  - *nomial* coefficients. These include: incidence coefficients such as binomial or  $q$ -Gaussian ones for finite geometries or fibonomial coefficients [4,5] which are not incidence coefficients. Here these  $F$  - *nomial* numbers are introduced also via  $c_2$ ) axiom in the Definition 1 from [1]:

$$(1) \quad |a \odot b| = \frac{f(a \odot b)}{f(a)f(b)} = \binom{n}{k}_F.$$

We notice with emphasis and not only occasionally, that  $c_2$ ) **axiom** in Definition 1 from [1] **is equivalent to the fundamental Theorem 1** from [1]. More than that - in Section 3 we shall see that all objects from Equation 1 gain specific uniform

combinatorial interpretation within the class of cobweb prefab combinatorial scheme  
- by construction.

## 2 Cobweb posets - presentation and their combinatorial interpretation

Given any sequence  $\{F_n\}_{n \geq 0}$  of nonzero reals one defines its corresponding binomial-like  $F$ -nomial coefficients in the spirit of Ward's Calculus of sequences [6] (reals may be replaced for example by any field of characteristic zero) as follows

**Definition 1**

$$\binom{n}{k}_F = \frac{F_n!}{F_k! F_{n-k}!} \equiv \frac{n_F^k}{k_F!}, \quad n_F \equiv F_n \neq 0, n \geq 0$$

where we make an analogy driven identifications in the spirit of Ward's Calculus of sequences ( $0_F \equiv 0$ ):

$$n_F! \equiv n_F(n-1)_F(n-2)_F(n-3)_F \dots 2_F 1_F;$$

$$0_F! = 1; \quad n_F^k = n_F(n-1)_F \dots (n-k+1)_F.$$

This is just the adaptation of the notation for the purpose Fibonomial Calculus case (see Example 2.1 in [7]). Given any such sequence  $\{F_n\}_{n \geq 0} \equiv \{n_F\}_{n \geq 0}$  of now

**nonzero integers** we define following [4,5] the partially ordered graded infinite set  $P$  - called afterwards a cobweb poset - as follows. Its vertices are labelled by pairs of coordinates:  $\langle i, j \rangle \in N \times N_0$  where  $N_0$  denotes the nonnegative integers. Vertices show up in layers ("generations") of  $N \times N_0$  grid along the recurrently emerging subsequent  $s$ -th levels  $\Phi_s$  where  $s \in N_0$  i.e.

**Definition 2**

$$\Phi_s = \{\langle j, s \rangle | 1 \leq j \leq s_F\}, s \in N_0.$$

We shall refer to  $\Phi_s$  as to the set of vertices at the  $s$ -th level. The population of the  $k$ -th level ("generation") counts  $k_F$  different member vertices for  $k > 0$  and one for  $k = 0$ . Here down a disposal of vertices on  $\Phi_k$  levels is visualized for the case of Fibonacci sequence (the subtlety of  $F_0 = 0$  is manageable)

```

--- ↑ --- ↑ --- up --- Fibonacci --- stairs --- * --- k-th-level

--- and --- so --- on --- up --- ↑ ---
*****10-th-level
*****9-th-level
*****8-th-level
*****7-th-level
*****6-th-level
*****5-th-level
***4-th-level
**3-rd-level
*2-nd-level
*1-st-level
*0-th-level

```

**Figure 1.** The  $s$ -th levels in  $N \times N_0$ ,  $N_0$  - nonnegative integers

Accompanying the set  $E$  of edges to the set  $V$  of vertices - we obtain the Hasse diagram where here down  $p, q, s \in N_0$ . (Convention: Edges stay for arrows directed - say - upwards) Namely:

**Definition 3**

$$P = \langle V, E \rangle, \quad V = \bigcup_{0 \leq p} \Phi_p, \quad E = \{ \langle \langle j, p \rangle, \langle q, (p+1) \rangle \rangle \} \cup \{ \langle \langle 1, 0 \rangle, \langle 1, 1 \rangle \rangle \},$$

where  $1 \leq j \leq p_F, 1 \leq q \leq (p+1)_F$ .

**Definition 4** The finite cobweb sub-poset  $P_m = \bigcup_{0 \leq s \leq m} \Phi_s$  is called the prime cobweb poset.

In reference [3,4] a partially ordered infinite set  $P$  was introduced via descriptive picture of its Hasse diagram. Indeed, we may picture out the partially ordered infinite set  $P$  from the Definition 3 with help of the sub-poset  $P_m$  (rooted at  $F_0$  level of the poset) to be continued then ad infinitum in now obvious way as seen from the figures Fig.1 – Fig.5 of  $P_m$  cobweb posets below. These look like the Fibonacci rabbits' way generated tree with a specific “cobweb” [4,5,8]. This is an example of acyclic directed graphs (DAG) [9] cobweb subclass.




Fig.1. Display of Natural numbers' cobweb poset.




Fig.2. Display of Even Natural numbers' cobweb poset.




Fig.4. Display of divisible by 3 natural numbers' cobweb poset.




Fig.5. Display of Fibonacci numbers' cobweb poset.

Compare with the bottom 6 levels of a Young-*Fibonacci lattice*, introduced by Richard Stanley- in Curtis Greene's gallery of posets: [www.haverford.edu/math/cgreene/posets/posetgallery.html](http://www.haverford.edu/math/cgreene/posets/posetgallery.html).

As seen above - for example the Fig.5. displays the rule of the construction of the Fibonacci "cobweb" poset. It is being visualized clearly while defining this cobweb poset  $P$  with help of its incidence matrix . The incidence  $\zeta$  function matrix representing uniquely just this cobweb poset  $P$  has the staircase structure correspondent with "cobwebbed" Fibonacci Tree i.e. a Hasse diagram of the particular partial order relation under consideration. This is seen below on the Fig.6 [8]:

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \dots \end{bmatrix}$$

Figure 6. The staircase structure of incidence matrix  $\zeta$  for the Fibonacci cobweb poset case

**Note** The knowledge of  $\zeta$  matrix explicit form enables one to construct (count) via standard algorithms [10] the Möbius matrix  $\mu = \zeta^{-1}$  and other typical elements of incidence algebra perfectly suitable for calculating number of chains, of maximal chains etc. in finite sub-posets of  $P$ . All elements of the corresponding incidence algebra are then given by a matrix of the Fig.5 with 1's replaced by any reals ( or ring elements in more general cases).

Right from the definition of  $P$  via its Hasse diagram here now follow quite obvious and important observations. They lead us to a combinatorial interpretation of cobweb poset's characteristic binomial-like coefficients (for example - fibonomial ones [4,5]). Here they are with the first obvious observation at the start.

**Observation 1**

*The number of maximal chains starting from The Root (level  $0_F$ ) to reach any point at the  $n$  - th level with  $n_F$  vertices is equal to  $n_F!$ .*

**Observation 2** ( $k > 0$ )

*The number of maximal chains **rooted in any** vertex at the  $k$  - th level reaching the  $n$  - th level with  $n_F$  vertices is equal to  $n_F^m$ , where  $m + k = n$ .*

Indeed. Denote the number of ways to get along maximal chains from **any point** in  $\Phi_k$  to  $\Rightarrow \Phi_n, n > k$  with the symbol

$$[\Phi_k \rightarrow \Phi_n]$$

then obviously we have :

$$[\Phi_0 \rightarrow \Phi_n] = n_F!$$

and

$$[\Phi_0 \rightarrow \Phi_k] \times [\Phi_k \rightarrow \Phi_n] = [\Phi_0 \rightarrow \Phi_n].$$

In order to formulate the combinatorial interpretation of  $F$ -sequence-nomial coefficients ( $F$ -nomial - in short) [6,4,5,8] let us consider all finite "max-disjoint" sub-posets rooted at the  $k$  - th level at any fixed vertex  $\langle r, k \rangle, 1 \leq r \leq k_F$  and ending at corresponding number of vertices at the  $n$  - th level ( $n = k + m$ ) where the "max-disjoint" sub-posets are defined below.

**Definition 5** *Two isomorphic copies of  $P_m$  are said to be max-disjoint if being considered as sets of maximal chains they are disjoint i.e they have no maximal chain in common. All of  $P_m$ 's constitute from now on a family of prime [1] prefabiants.*

**Definition 6** *We denote the number of all max-disjoint isomorphic copies of  $P_m$  rooted at any vertex  $\langle j, k \rangle, 1 \leq j \leq k_F$  of  $k$  - th level with the symbol*

$$\binom{n}{k}_F.$$

We use the accustomed to practical convention:  $\binom{0}{0}_F = 1$ .

Naturally the above definition make sense not for arbitrary  $F$  sequences as  $F$ -nomial coefficients should be nonnegative integers.

**Definition 7** *A sequence  $F = \{n_F\}_{n \geq 0}$  is called cobweb-admissible iff*

$$\binom{n}{k}_F \in N \cup \{0\} \quad \text{for} \quad k, n \in N \cup \{0\} \equiv Z_{\geq}.$$

Recall now that the number of ways to reach an upper level from a lower one along any of maximal chains i.e. the number of all maximal chains from the level  $\Phi_k$  to  $\Rightarrow \Phi_n$ ,  $n > k$  is equal to

$$[\Phi_k \rightarrow \Phi_n] = n_F^m.$$

Naturally then we have

$$(2) \quad \binom{n}{k}_F \times [\Phi_0 \rightarrow \Phi_m] = [\Phi_k \rightarrow \Phi_n] = n_F^m$$

where  $[\Phi_0 \rightarrow \Phi_m] = m_F!$  counts the number of maximal chains in any copy of the  $P_m$ . With this in mind we see that the following holds.

**Observation 3** ( $n, k \geq 0$ )

Let  $n = k + m$ . The number of max-disjoint sub-posets isomorphic to  $P_m$  (max-disjoint isomorphic copies of prime prefabiants), rooted at the  $k$ -th level and ending at the  $n$ -th level is equal to

$$\begin{aligned} \frac{n_F^m}{m_F!} &= \binom{n}{m}_F \\ &= \binom{n}{k}_F = \frac{n_F^k}{k_F!}. \end{aligned}$$

**Note** The Observation 3 provides us with the **new combinatorial interpretation** of the class of all classical  $F$ -nomial coefficients including distinguished binomial or distinguished Gauss  $q$ -binomial ones or Konvalina generalized binomial coefficients of the first and of the second kind [11]- which include Stirling numbers too. The vast family of Ward-like [6] admissible by  $\psi = \langle \frac{1}{n_F!} \rangle_{n \geq 0}$ -extensions  $F$ -sequences [7,12] includes also those desired here which shall be called "GCD-morphic" sequences. This means that  $GCD[F_n, F_m] = F_{GCD[n,m]}$  where  $GCD$  stays for Greatest Common Divisor operator. The Fibonacci sequence is a much nontrivial and guiding famous example of GCD-morphic sequence. Naturally **incidence coefficients of any reduced incidence algebra of full binomial type** [13] are GCD-morphic sequences therefore they are now independently given a new cobweb combinatorial interpretation via Observation 3. More on that - see the next section where prefab combinatorial description is being served. Before that - on the way - let us formulate the following problem (open?).

**Problem 1** Find effective characterizations of the cobweb admissible sequence i.e. find all examples.

### 3 Cobweb posets as prefabs with *nonassociative* and noncommutative composition

Finite cobweb sub-posets i.e. isomorphic copies of  $P_m, m \geq 0$  constitute connected acyclic digraphs as well as the Hasse diagram of the infinite cobweb poset  $P$  is. Directed acyclic graphs are denoted as DAG's [9]. Hence one might call connected DAG's - directed trees. As for the recent development on acyclic digraphs we refer to [14] and references therein. As in [14] one considers here digraphs on labeled vertices and a "digraph" means a simple graph with at most one edge directed from vertex to vertex. Loops and cycles of length two are permitted in general, but parallel edges are forbidden. "Acyclic" means that there are no cycles of any length. Apart from Theorem 1 there note in [14] also Bibliographic remarks on acyclic digraphs refereing

to Robinson and Stanley and then to Bender et al. and Gessel. Because of an easy access to Plotnikov's paper [9] we shall take other definitions from there - if needed - for granted. These are temporarily used just for the guiding observation relating cobweb prefabs' digraphs to [9]. Namely - in terminology of [9] - we make rather obvious observation.

**Observation 4** *The Hasse (here upward oriented) diagram of any prime cobweb poset or  $P$  is an oDAG.*

For the sake of explanation we quote after [9]: A poset  $P$  is of the *dimension* 2;  $\dim P = 2$  if there exist two chains  $L_1$  and  $L_2$  such that  $P = L_1 \cap L_2$ .

A digraph  $G$  is called *the orderable digraph (oDAG)* if there exists a  $\dim 2$  poset such that its Hasse diagram coincides with the digraph  $G$ .

We shall pass over now to the brief presentations of a cobweb prefab combinatorial structure [1] in which each object (*prefabiant*) is uniquely representable by construction as a synthesis (composition) of powers of prime objects where here these are the cobweb sub-posets  $P_n$  of  $P$  that are to be identified with prime *prefabiants*. Since now on we shall adhere to the notation and terminology of [1]. We assume the acquaintance of [1] which is justly considered as famous as important.

The definition of prefab combinatorial structure  $(S, \odot, f)$  here is assumed to be given by Definition 1 from [1] except for associativity requirement  $a_1$ ) and commutativity requirement  $a_2$ ), which are postponed until stated otherwise. In general  $a \odot b \neq b \odot a, a, b \in S$  - already for prime objects. The definition of weighted (not necessarily associative, commutative) prefab and enumerator  $g(A), A \subseteq S$  are then Definitions 2 and 3 from [1] correspondingly. We shall now formulate Observation 5 (to to be checked by careful examination)- observation of distinguished importance for the combinatorial interpretation of the **property  $c_2$ ) from the Definition 1** [1] of the prefab. The property  $c_2$ ) postulate from [1] is

$$(3) \quad |a \odot b| = \frac{f(a \odot b)}{f(a)f(b)}, \quad a, b \in S,$$

whenever  $a$  and  $b$  have no common factor different from identity prefabiant  $i$  [1] where here  $|A|$  denotes here the number of max-disjoint isomorphic copies of prime prefabiants in the set  $A = a \odot b$ . The function  $f$  satisfies the requirement  $c_1$  - of course.

**Observation 5** *Let the enumerator or generating function for prefab subsets be defined as indicated above. Then the set of requirements  $Prefab_{c(2)} = \{a_3), b_1), b_2), c_1), c_2)\}$  is equivalent to set of requirements  $Prefab_{(Th.1)} = \{a_3), b_1), b_2), c_1), Theorem.1\}$ , where Theorem 1 means Theorem 1 from [1].*

Both sets of requirements define on  $S$  the same prefab structure (not necessarily commutative and associative) where requirements  $b_1), b_2)$  are to be understood as rewritten in an order and brackets being taken into account fashion.

Now comes the example of the class of weighted prefabs  $(S, \odot, f, \omega)$  with noncommutative, *nonassociative* synthesis (composition)  $\odot$ . We shall call this binary multivalued operation' [1] analogue case here a "*coopt-synthesis*"  $\odot$ .

**Cobweb prefab combinatorial structure.** The family  $S$  of combinatorial objects (*prefabiants*) consists of all layers  $\langle \Phi_k \rightarrow \Phi_n \rangle, k < n, k, n \in N \cup \{0\} \equiv Z_{\geq}$  and an empty prefabiant  $i$ . Layer is considered here to be the set of all max-disjoint isomorphic copies (iso-copies) of  $P_{n-k}$ . The set  $\wp$  of prime objects consists of all sub-posets  $\langle \Phi_0 \rightarrow \Phi_m \rangle$  i.e. all  $P_m$ 's  $m \in N \cup \{0\} \equiv Z_{\geq}$  constitute from now on a family of prime *prefabiants* [1]. The  $Z_{\geq}$  grading preserving  $\odot$  coopt-synthesis for prime prefabiants  $P_k \odot P_m = \langle \Phi_k \rightarrow \Phi_n \rangle, n = k + m$  means: consider the leafs of  $P_k$  to be the roots of all max-disjoint isomorphic copies (iso-copies) of  $P_m$ . Run through all the leafs (now - roots). The  $Z_{\geq}$  grading preserving  $\odot$  synthesis of (not necessarily prime)



prefabiants - accordingly means the same procedure with the requirement added (see: Example 5 in [1]). If this algorithm applied to subsequent prime elements of the second prefab gives rise to a layer of max-disjoint prefabs more then one way - keep only one copy of it. As a result we have:

$$\langle \Phi_k \rightarrow \Phi_n \rangle \odot P_s = \langle \Phi_n \rightarrow \Phi_{n+s} \rangle, \quad k \in Z_{\geq}, s \in N, n > k.$$

Accordingly the  $Z_{\geq} \times Z_{\geq}$  grading of  $S$  preserving  $\odot$  synthesis ( $\odot$  *coopt-synthesis*) is defined for arbitrary elements of  $S$  as simply as follows:

$$\langle \Phi_k \rightarrow \Phi_n \rangle \odot \langle \Phi_t \rightarrow \Phi_{t+s} \rangle = \langle \Phi_n \rightarrow \Phi_{n+s} \rangle; \quad t, k \in Z_{\geq}, n > k, s > 0.$$

In order to satisfy the requirement  $a_3$ ) we postulate for an empty prefabiant  $i$  that

$$\langle \Phi_k \rightarrow \Phi_n \rangle \odot i = i \odot \langle \Phi_k \rightarrow \Phi_n \rangle = \langle \Phi_k \rightarrow \Phi_n \rangle, \quad k \in Z_{\geq}, n > k.$$

The appropriately adjusted requirements  $b_1), b_2)$  are satisfied by construction as

$$\langle \Phi_k \rightarrow \Phi_n \rangle = P_k \odot P_{n-k}, \quad k \in Z_{\geq}, n > k.$$

The coopt-synthesis  $\odot$  is nonassociative by construction as

$$(\langle \Phi_k \rightarrow \Phi_n \rangle \odot \langle \Phi_t \rightarrow \Phi_{t+s} \rangle) \odot \langle \Phi_p \rightarrow \Phi_{p+q} \rangle = P_{n+s} \odot P_q; \quad p, t, k \in Z_{\geq}, n > k, s > 0, q > 0$$

while

$$\langle \Phi_k \rightarrow \Phi_n \rangle \odot (\langle \Phi_t \rightarrow \Phi_{t+s} \rangle \odot \langle \Phi_p \rightarrow \Phi_{p+q} \rangle) = P_n \odot P_q; \quad p, t, k \in Z_{\geq}, n > k, s > 0, q > 0.$$

As for the size functions let us aid with an analogy (see: Example 5 in [1]).

**Analogy:**

*Graphs.....Cobweb – prefabs    I.....Cobweb – prefabs    II*

*vertices.....max – chains.....leafs    of     $P_n$ 's*

*connected    $G_n$    on    $[n]$    ..... $P_n$    cobweb..... $P_n$    cobweb*

*size( $G_n$ ) =  $n$ .....size<sub>1</sub>( $P_n$ ) =  $n!$ .....size<sub>2</sub>( $P_n$ ) =  $n$*

*$f(G_n) = n!$ ..... $f(P_n) = n!$ ..... $f(P_n) = n!$    .*

Recall now the Equation 3. We may draw now from all the above the following **conclusion**.

#### **Conclusion I**

In the finite cobweb posets setting the  $f$  function may be chosen so as to be the *size<sub>1</sub>* of a prime prefabiant with  $n_F$  leafs or so as to be factorial of the *size<sub>2</sub>* of a prime prefab with  $n_F$  leafs. This gives:

$$f(P_n) = n_F!, \quad f(P_n^k) = (km)_F!$$

and the Equation 1 gets the required, expected combinatorial interpretation for any cobweb prefab structure determined by the choice of any sequence of natural numbers from the countless family of cobweb admissible sequences. Thus we are equipped with the cobweb prefab's uniform combinatorial interpretation of all of them at once.

### Conclusion II

Naturally the Corollary 1 from [1] also holds in our case. Choosing now the weight function to be of the form

$$\omega(a) = x^n, \quad n = \text{size}_2(a), \quad a \in S$$

we have the weighted cobweb prefab and consequently (see: Examples 5,10 in [1]) the formula for the cobweb weighted prefab enumerator reads

$$(4) \quad g(S) = \exp\{g(\varphi)\},$$

where

$$(5) \quad g(\varphi) = \exp_F\{x\} - 1,$$

while

$$(6) \quad \exp_F\{x\} = \sum_{n \geq 0} \frac{x^n}{n_F!}$$

$\exp_F$  function [6,12,7,15] - is the primary object of extended finite operator calculus being recently developed in [7,12,15]. There it serves to define a central object of extended umbral calculus i.e. the generalized translation operator  $E^a(\partial_F)$  where the linear difference operator  $\partial_F$ ;  $\partial_F x^n = n_F x^{n-1}$ ;  $n \geq 0$  is known under the name of the  $F$ -derivative [6,7,14,15]. here comes the example (11) from [1] interpreted in the language of  $\psi$  - extensions [6] in their operator form [12,15,7].

**Bender - Goldman - prefab example** Let the "prefabian"  $\hat{q}$ -Bell numbers  $B_n^{pref}(\gamma)$  be defined as sums over  $k$  of  $\hat{S}_q(n, k)$  Stirling numbers equal of the number of unordered direct sums decompositions of the  $n$ -dimensional vector space  $V_{q,n}$  over  $GF(q) \equiv F_q$  with  $k$  summands. Then the Bender-Goldman exponential formula (17) from [1] in  $\psi$ -extensions' notation [12,15,7] reads

$$B_\gamma^{pref}(x) = \sum_{n \geq 0} B_n^{pref}(\gamma) \frac{x^n}{n_\gamma!} = \exp\{\exp_\gamma(x) - 1\}. \quad (\gamma - e.g.f.)$$

Here

$$n_\gamma! = (q^n - 1)(q^n - q^1) \dots (q^n - q^{n-1}) = |GL_n(F_q)|,$$

$D_o(q) = 1$  by convention while  $D_n(q) \equiv B_n^{pref}(\gamma) =$  number of all unordered direct sums decompositions of the vector space  $V_{q,n}$ .

### The natural hint

The appealing analogy of the above schema and example just presented give rise to questions on their eventual correspondents as  $Stirling_F$  numbers of the second kind,  $\exp_F$ -ponential polynomials and  $F$ -Dobinski like formulas. Such extensions are more or less implicit in some papers . for example - see Wagner's (1.15) formula in [16] which formally becomes of the  $(\gamma - e.g.f.)$  formula form from above with now almost arbitrary  $\gamma = \langle \frac{1}{n_\gamma!} \rangle_{n \geq 0}$  sequence (see also [17] and references therein). These questions are to be considered elsewhere. As for the related (determined by  $F - nomial$ 's) extended umbral calculi in its operator form one may contact also very recent review [18].

## 4 Cobweb posets as prefabs with associative and commutative composition

Here another single valued commutative and associative composition case is presented in brief. The definition of the next prefab combinatorial structure with the single valued composition  $(S, \circ, f)$  is assumed to be given here by the Definition 1 from [1] including associativity requirement  $a_1$ ) and commutativity requirement  $a_2$ ).

The family  $S$  of combinatorial objects (*prefabiants*) consists now of all layers  $\langle \Phi_k \rightarrow \Phi_n \rangle$ ,  $k < n$ ,  $k, n \in N \cup \{0\} \equiv Z_{\geq}$  and an empty prefabiant  $i$  to be interpreted as the name or representant of all "empty layers"  $\langle \Phi_m \rightarrow \Phi_m \rangle$ . Layer is considered here as the set of all max-disjoint isomorphic copies (iso-copies) of  $P_{n-k}$ . The set  $\varnothing$  of prime objects consists of all sub-posets  $\langle \Phi_0 \rightarrow \Phi_m \rangle$  i.e. all  $P_m$ 's  $m \in N \cup \{0\} \equiv Z_{\geq}$  constitute from now on a family of prime *prefabiants* [1].

The  $Z_{\geq}$  grading preserving  $\circ$  coopt-synthesis for prime *prefabiants*  $P_k \circ P_m = \langle \Phi_0 \rightarrow \Phi_{n+m} \rangle$ ,  $n = k+m$  means: consider the leafs of  $P_k$  to be the *transitory* roots of all max-disjoint isomorphic copies (iso-copies) of  $P_m$ . Run through all the leafs (now -transitory roots).

The  $Z_{\geq} \times Z_{\geq}$  grading of  $S$  grading preserving  $\circ$  synthesis of (not necessarily prime) prefabiants - accordingly means the same procedure with the requirement added (see: Example 5 in [1]): if this algorithm applied to subsequent prime elements of the second prefab gives rise to a layer of max-disjoint prefabs more then one way - keep only one copy of it. As a result we have:

$$\langle \Phi_k \rightarrow \Phi_n \rangle \circ P_s = \langle \Phi_{k+0} \rightarrow \Phi_{n+s} \rangle, \quad k \in Z_{\geq}, s \in N, n > k.$$

Accordingly the  $Z_{\geq} \times Z_{\geq}$  grading of  $S$  preserving *coopt-synthesis* " $\circ$ " is defined as follows:

$$\langle \Phi_k \rightarrow \Phi_n \rangle \circ \langle \Phi_p \rightarrow \Phi_q \rangle = \langle \Phi_{k+p} \rightarrow \Phi_{n+q} \rangle; \quad p, k \in Z_{\geq}, n > k, q > p.$$

### Conclusion III

In this finite cobweb posets setting with associative and commutative composition  $\circ$  being single valued the  $f$  function may be chosen as constant equal to 1 function (note the other possibilities:  $f(\langle \Phi_k \rightarrow \Phi_n \rangle) = \alpha^{n-k}$ ,  $\alpha \neq 0, n > k$ ).

This implies validity of the Corollary 2 and the Corollary 3 from [1] also in the cobweb " $\circ$  - case" - providing us with direct efficient analogy to the cases of unlabeled graphs, unordered partitions or factorizations of integers (see Examples 1,2,3 in [1]).

### Acknowledgements

Discussions with Participants of Gian-Carlo Rota Polish Seminar  
<http://ii.uwb.edu.pl/akk/index.html> - are highly appreciated.

## References

- [1] E. Bender, J. Goldman *Enumerative uses of generating functions*, Indiana Univ. Math.J. **20** 1971), 753-765.
- [2] D. Foata and M. Sch"utzenberger, Th'eorie g'eometrique des polynomes euleriens, (Lecture Notes in Math., No. 138). Springer-Verlag, Berlin and New York, 1970.
- [3] A. Nijenhuis and H. S. Wilf, *Combinatorial Algorithms*, 2nd ed., Academic Press, New York, 1978.
- [4] A. K. Kwasniewski *Information on combinatorial interpretation of Fibonomial coefficients* Bull. Soc. Sci. Lett. Lodz Ser. Rech. Deform. 53, Ser. Rech.Deform. **42** (2003), 39-41. ArXiv: math.CO/0402291 v1 18 Feb 2004

- [5] A. K. Kwaśniewski, *The logarithmic Fib-binomial formula* Advanced Stud. Contemp. Math. **9** No 1 (2004), 19-26
- [6] M. Ward: *A calculus of sequences*, Amer.J.Math. Vol.58, (1936), 255-266.
- [7] A. K. Kwaśniewski, *On simple characterizations of Sheffer  $\Psi$ -polynomials and related propositions of the calculus of sequences*, Bull. Soc. Sci. Lettres Łódź **52**, Sér. Rech. Déform. **36** (2002), 45-65. ArXiv: math.CO/0312397 2003
- [8] A. K. Kwaśniewski, *More on combinatorial interpretation of fibonomial coefficients*, Bull. Soc. Sci. Lettres Łódź **54**, Sér. Rech. Déform. **44** (2004), 23-38 –65. ArXiv: math.CO/0402344 v1 22 Feb 2004
- [9] A.D. Plotnikov *A formal approach to the oDAG/POSET problem* (2004) [html://www.cumulativeinquiry.com/Problems/solut2.pdf](http://www.cumulativeinquiry.com/Problems/solut2.pdf) (submitted to publication - March 2005)
- [10] E.Krot: *The first ascent into the Fibonacci Cob-web Poset* (submitted to publication - December 2004), ArXiv: math.CO/0411007
- [11] J. Konvalina, *A Unified Interpretation of the Binomial Coefficients, the Stirling Numbers and the Gaussian Coefficients* The American Mathematical Monthly **107**(2000), 901-910.
- [12] A. K. Kwaśniewski *Main theorems of extended finite operator calculus* Integral Transforms and Special Functions, **14** No 6 (2003), 499-516.
- [13] E. Spiegel, Ch. J. O'Donnell *Incidence algebras* Marcel Dekker, Inc. Basel 1997.
- [14] Brendan D. McKay, Frederique E. Oggier, Gordon F. Royle, N. J. A. Sloane, Ian M. Wanless and Herbert S. Wilf, *Acyclic digraphs and eigenvalues of (0,1)-matrices* Journal of Integer Sequences, **7**, August 2004 Article 04.3.3 (arXiv: math.CO/0310423)
- [15] A. K. Kwaśniewski: *On Extended Finite Operator Calculus of Rota and Quantum Groups*, Integral Transforms and Special Functions Vol 2, No 4, (2001), 333-340
- [16] Carl G. Wagner *Generalized Stirling and Lah numbers* Discrete Mathematics **160** (1996), 199-218.
- [17] A. K. Kwaśniewski: *Information on some recent applications of umbral extensions to discrete mathematics* to appear in Review Bulletin of Calcutta Mathematical Society Vol 13 ( 2005) ArXiv: math.CO/0411145 7 Nov 2004
- [18] A.K. Kwaśniewski, E. Borak: *Extended finite operator calculus - an example of algebraization of analysis* Central European Journal of Mathematics **2** (5), 2005, 767-792. ArXiv math.CO/0412233 14 Dec 2004